



Sugarcane Crop Nutrition Schedule - North India

Ratoon - cane



GROWTH STAGE

Fertiliser application	Basal application	30-45 days after ratooning	60-75 days after ratooning	90-105 days after ratooning	120-250 days after ratooning
Polysulphate	100 kg/acre			25 kg/acre	
FertiFlow 5-45-5 + 8Zn		5 kg/acre			
FertiFlow 12-6-22 + 12CaO			5 kg/acre	5 kg/acre	
UREA (46% N)	45 kg/acre	45 kg/acre	45 kg/acre	45 kg/acre	
DAP (18% N, 46% P ₂ O ₅)	50 kg/acre				
MOP (60% K ₂ O)				50 kg/acre	
NutriVant - Starter (11-36-24 + TE)		10 g/l water			
Nutrivant - PeakVant (00-49-32)			10 g/l water		
Nutrivant - Booster (8-16-39 + TE)				10 g/l water	
Boron (Foliar grade)		1 g/l water		1 g/l water	
Fe - EDTA (12%)			2 g/l water		

The recommendations are for general information purpose only and should not be considered as advice or consultancy. The exact fertilisation program should be determined by the customer according to the crop selection, crop yield, soil and water conditions and the growing conditions, which are prone to changes due to external factors from time to time. The recommendations made are generic in nature. Please consult a local agronomist in advance for implementation and advice, specifically.

- Soil application
- Foliar application
- Side dressing



Polysulphate FertiFlow™ **NutriVant™**
Foliar Nutrition Line

Toll free number
1800 210 3031